



SUNBURN

Sunburn is most commonly seen on the trunk of affected trees, where direct exposure to the sun and high temperatures can cause damage to the cambium layer. This presents as cracking, splitting, peeling, or discolored bark.


WHAT TO LOOK OUT FOR:

Sunburn damage will occur during the summer but will remain year round. It can be found on the southwest side of the trunk and on the upper side of limbs.



SIGNS OF SUNBURN:

Sunburn will cause discolored patches to form on areas of bark that receive direct light. In severe cases, the bark will crack or split entirely and ooze sap.



HOW TO TREAT:

PREVENTION

Young trees are especially prone to sunburn until their canopies grow wide enough to shade the trunk. Prevent any potential damage by applying trunk wrap or paint on the trunk of the tree, from the base to the first main branch of the canopy. Any limbs receiving direct exposure can be protected, too.

FERTILIZER

Sunburn on the trunk of the tree can cause stress throughout the entire plant. Combat the stress and give your trees all the nutrients they need to heal from the damage with applications of Super Charged Moon Juice™ and Moon Microboost™.



SCAN HERE FOR MORE INFORMATION!

SUNBURN TREATMENT PACKAGE: **\$119⁹⁹**



SUPER CHARGED MOON JUICE™

+



MOON MICROBOOST™

+



TREE WRAP

+



TRUNK PAINT

Recommended products are not guaranteed to resolve issue. Always follow all instructions on chemical labels. Call or see store for questions.